

# Flux-mediated Dark Matter

Ji-Seon Song

Chung-Ang University


JHEP 06(2021) 013

In collaboration with Yoo-Jin Kang, Hyun Min Lee, Adriana G. Menkara

# Outline

- Hierarchy problems & Selfish Higgs
- Flux-mediated Dark Matter
- Result

# Hierarchy Problems



## Higgs Mass

- In SM, all physical quantities are represented by VEV of the Higgs field
- VEV of the Higgs field is  $10^{15}$  times smaller than the Planck mass  
→ Why the Higgs boson is so much lighter than the Planck mass?


## Cosmological Constant

- Theoretically, the value of the cosmological constant can be approximately predicted as a Planck mass
- But measured value of the cosmological constant is  $10^{120}$  times smaller than the Planck mass  
→ Why is it so small?

# Selfish Higgs

- Dynamical relaxation of Higgs mass and Cosmological constant
- Higgs acts as an anthropic selector for the emergence of a fairly unique non-empty universe

→ called **Selfish Higgs**



- Selfish Higgs is based on the SM with addition of a single non-dynamical field described by a four-form


# Four-Form Flux

- The four-form derives from a three-form

$A_{\nu\rho\sigma}$  : Three-index anti-symmetric tensor field

Four-form field strength :  $F_{\mu\nu\rho\sigma} = 4\partial_{[\mu}A_{\nu\rho\sigma]}$

Giudice et al, Kaloper et al. 2019





- Assume that the four-form starts with a large flux parameter in the early universe in a highly excited state
  - excited **four-form** can reduce its energy by **creating membranes**
  - four-form undergoes a configuration transition ( $n \rightarrow n-1$ )
- After last membrane nucleation, the universe would be empty **without reheating**

# Model for four-form and Dark Matter

We consider **pseudo-scalar field** ( $\phi$ ) for reheating & **Dirac fermion dark matter** ( $\chi'$ )

➤  $\mathcal{L} = \mathcal{L}_0 + \mathcal{L}_{relax} + \mathcal{L}_{RH} + \mathcal{L}_{DM} + \mathcal{L}_S + \mathcal{L}_L + \mathcal{L}_{memb}$


$$\mathcal{L}_0 = \sqrt{-g} \left[ \frac{1}{2}R - \Lambda - \frac{1}{48}F_{\mu\nu\rho\sigma}F^{\mu\nu\rho\sigma} - |D_\mu H|^2 - M^2|H|^2 + \lambda_H|H|^4 \right]$$
$$\mathcal{L}_{relax} = \frac{c_H}{24}\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu\rho\sigma}|H|^2 \quad \text{Higgs mass scanning}$$
$$\mathcal{L}_{RH} = -\frac{1}{2}(\partial_\mu\phi)^2 - \frac{1}{2}m_\phi^2(\phi - \alpha)^2 + \frac{\mu}{24}\epsilon^{\mu\nu\rho\sigma}F_{\mu\nu\rho\sigma}\phi \quad \text{reheating}$$
$$\mathcal{L}_{DM} = i\bar{\chi}'\gamma^\mu\partial_\mu\chi' - m'_\chi\bar{\chi}'\chi' + i\frac{m'_\chi}{f}\phi\bar{\chi}'\gamma^5\chi' \quad \text{Dark matter}$$



# The Relaxation Mechanism with Four-Form Flux

- Using equation of motion for  $F_{\mu\nu\rho\sigma}$

$$\rightarrow \mathcal{L} = \sqrt{-g} \left[ \frac{1}{2}R - \Lambda - |D_\mu H|^2 + M^2|H|^2 - \lambda_H|H|^4 \right] \\ - \frac{1}{2}(\partial_\mu \phi)^2 - \frac{1}{2}m_\phi^2(\phi - \alpha)^2 - \frac{1}{2}(\mu\phi + c_H|H|^2 + q)^2 + \mathcal{L}_{DM} + \mathcal{L}_{nucl}$$

- Equation of motion for  $A_{\nu\rho\sigma}$  :  $\epsilon^{\mu\nu\rho\sigma}\partial_\mu q = -e \int d^3\xi \delta^4(x - x(\xi)) \frac{\partial x^\nu}{\partial \xi^a} \frac{\partial x^\rho}{\partial \xi^b} \frac{\partial x^\sigma}{\partial \xi^c} \epsilon^{abc}$   
→ Flux parameter  $q$  is quantized in units of  $e$  as  $q = en$  ( $n$  : integer)
- Nucleate a membrane :  $q = en \rightarrow e(n-1) \rightarrow e(n-2) \rightarrow \dots$
- Flux change stops at  $q = q_c - e$  with  $q_c \equiv M^2/c_H - \mu\langle\phi\rangle$

$$\rightarrow \left[ \begin{array}{l} M_{eff}^2(q) = M^2 - c_H(q + \mu\langle\phi\rangle) \\ \Lambda_{eff}(q) = \Lambda + \frac{1}{2}q^2 + V(\langle\phi\rangle) + V(\langle H \rangle) \end{array} \right. \quad \begin{array}{l} \xrightarrow{} c_H e \sim (100 GeV)^2 \\ \xrightarrow{} \sim 0 \end{array} \quad \left. \right\} \text{relaxed}$$

# Four-form coupling induced Higgs mixing

- SM Higgs & pseudo-scalar are expanded around the vacuum :

$$\langle H \rangle = \frac{(0, v_H(q) + h)^T}{\sqrt{2}} \quad \& \quad \langle \phi \rangle = v_\phi + \varphi$$

$$v_H(q) = \sqrt{\left(M^2 - c_H(q + \mu v_\phi)\right) / \left(\lambda_H + \frac{1}{2}c_H^2\right)}$$

$$v_\phi(q) = \frac{m_\phi^2}{\mu^2 + m_\phi^2} \left[ \alpha - \frac{\mu}{m_\phi^2} \left( \frac{1}{2} c_H v_H^2 + q \right) \right]$$

- Transformation to the mass eigenstates  $(h_1, h_2)$

$$\rightarrow m_{h_{1,2}}^2 = \frac{1}{2} (m_\phi^2 + m_h^2) \mp \frac{1}{2} \sqrt{(m_\phi^2 - m_h^2)^2 + 4c_H^2 \mu^2 v_H^2(q)} \quad \& \quad \tan 2\theta(q) = \frac{2c_H \mu v_H(q)}{m_\phi^2 - m_h^2}$$

- Electroweak symmetry is broken at  $q = q_c - e$

$$v^2 = \frac{m_\phi^2}{\mu^2 + m_\phi^2} \left( \frac{c_H e}{\lambda_{H,eff} - \frac{1}{2} \frac{c_H^2 \mu^2}{\mu^2 + m_\phi^2}} \right)$$

# Reheating

- Just after the last membrane nucleation, **pseudo-scalar VEV is shifted** by four-form flux

$$\Delta v_\phi = v_{\phi,c} - v_{\phi,0} = \frac{\lambda_H}{c_H} \frac{v^2 \mu}{m_\phi^2}$$

- The initial vacuum energy for reheating :  $V_i = \frac{1}{2} \left( \frac{\lambda_H}{c_H} \frac{\mu^2}{m_\phi^2} - \frac{1}{2} c_H \right)^2 v^4$


→ The **maximum temperature** of the Universe after inflation :

$$T_{\max} \cong 40 \text{GeV} \left( \frac{V_i^{1/4}}{100 \text{GeV}} \right) \left( \frac{100}{g_*} \right)^{1/4}$$

- Choose  $\sqrt{e} \sim 100 \text{GeV} - 10 \text{TeV}$  for  $\frac{m_\phi}{|\mu|} \sim 0.01 - 1$

$$\rightarrow T_{\max} \sim 40 \text{GeV} - 4 \text{TeV}$$

- For  $m_\chi < T_{\max}$ , DM starts being relativistic and in equilibrium



# Dark Matter Interactions

- Mass term for dark matter(DM) shifted (  $\because$  VEV of the pseudo-scalar field )

$$\mathcal{L}_{\chi, \text{mass}} = -m'_\chi \bar{\chi}' \chi' + \frac{i m'_\chi v_\phi}{f} \bar{\chi}' \gamma^5 \chi'$$

where  $m_\chi = m'_\chi \sqrt{1 + \frac{v_\phi^2}{f^2}}$  &  $\chi = (\chi_1, \chi'_2)^T$

$$= -m_\chi \bar{\chi} \chi$$

with  $\chi_1 = e^{-i\beta} \chi'_1$  &  $\tan\beta = \frac{v_\phi}{f}$

Redefined DM

- Interaction terms for dark matter :

$$\mathcal{L}_{\chi, \text{int}} = \frac{i m_\chi}{f} e^{-i\beta} (cos\theta(q) h_1 + sin\theta(q) h_2) \bar{\chi} P_R \chi - \frac{i m_\chi}{f} e^{-i\beta} (cos\theta(q) h_1 + sin\theta(q) h_2) \bar{\chi} P_L \chi$$

$$\equiv - \sum_{i=1,2} h_i \bar{\chi} (v_{\chi,i} + i a_{\chi,i} \gamma^5) \chi$$

$v_{\chi,1} = \frac{m_\chi}{f} sin\beta cos\theta(q)$  ,  $a_{\chi,1} = -\frac{m_\chi}{f} cos\beta cos\theta(q)$   
 $v_{\chi,2} = \frac{m_\chi}{f} sin\beta sin\theta(q)$  ,  $a_{\chi,2} = -\frac{m_\chi}{f} cos\beta sin\theta(q)$

- Yukawa couplings between the SM Higgs & SM fermions f :

$$\mathcal{L}_Y = -\frac{m_f}{v} h \bar{f} f$$

$$\equiv - \sum_{i=1,2} v_{f,i} h_i \bar{f} f$$


$v_{f,1} = -\frac{m_f}{v} sin\theta(q)$   
 $v_{f,2} = \frac{m_f}{v} cos\theta(q)$

# Dark Matter Interactions

- CP Violation


  - Four-form coupling to pseudo scalar field : **CP-odd**

  - Four-form coupling to Higgs field : CP-even



- Mixing induced by four-form flux

  - leads DM  $\rightarrow$  SM annihilation




- DM annihilation

  - Unsuppressed annihilation exists

- Direct Detection  $\rightarrow$  Suppressed

# DM annihilation


- We consider  $m_\chi \leq T_{max}$
- DM can pair annihilate into a pair of the SM fermions ( $\chi\bar{\chi} \rightarrow f\bar{f}$ )



$$(\sigma v_{rel})_{\chi\bar{\chi} \rightarrow f\bar{f}} = \frac{m_f^2 m_\chi^4}{8\pi v^2 f^2} \cos^4 \beta \sin^2 2\theta$$

$$\times \left( \frac{1}{4m_\chi^2 - m_{h_1}^2} - \frac{1}{4m_\chi^2 - m_{h_2}^2} \right)^2 \left( 1 - \frac{m_f^2}{m_\chi^2} \right)^{3/2}$$

- For  $m_\chi \geq m_{h_1}, m_{h_2}$



$\chi\bar{\chi} \rightarrow h_1 h_1$  : s-wave  $\rightarrow$  suppressed

p-wave  $\rightarrow$  unsuppressed

$\chi\bar{\chi} \rightarrow h_2 h_2$  : suppressed


$\chi\bar{\chi} \rightarrow h_1 h_2$  : suppressed

By Higgs mixing angle

$\theta \ll 1$

# DM Direct Detection


- $\mathcal{L}_{DD} = \sum_f \lambda_f \sum_{i=1,2} \frac{\tilde{v}_i}{m_{h_i}^2} [\bar{\chi}(v_{\chi,i} + i a_{\chi,i} \gamma^5) \chi] \bar{f} f$
- Elastic scattering between DM & nucleons



$$\sigma_{\chi-N} \simeq \frac{\mu_{\chi N}^2 m_\chi^2}{4\pi v^2 f^2 A^2} (sin2\theta)^2 \sin^2 \beta \left( \frac{1}{m_{h_1}^2} - \frac{1}{m_{h_2}^2} \right)^2 (Z f_p + (A-Z) f_n)^2$$

: doubly suppressed

- Elastic scattering between DM & electron





$$\sigma_{\chi-e} \simeq \frac{\mu_{\chi e}^2 m_e^2 m_\chi^2}{4\pi v^2 f^2} (sin2\theta)^2 \sin^2 \beta \left( \frac{1}{m_{h_1}^2} - \frac{1}{m_{h_2}^2} \right)^2$$



: doubly suppressed

# DM Direct Detection

- 1 – loop diagrams



$$M \sim v_\chi^2 v_q^2 \sin\theta^2 \frac{m_\chi^2}{m_\phi^4}$$





**Important for DD**  
 for sizable mixing and/or light  
 pseudo-scalar  
 (mixing between CP-odd & CP-even)

Work in progress

cf. Tomohiro Abe et al. 2019  
 (no CP violation)

# Results




- Red line : Relic density
  - Gray & Green : excluded by XENON1T and LEP , respectively
  - Purple : disfavored by diffuse gamma-rays from Fermi-LAT dwarf galaxies
  - Blue : favored by AMS-02 anti-proton excess
- DM & Nucleon scattering


$h_1 \rightarrow b\bar{b}$

# Results

$$m_\chi = 50 \text{ GeV}, m_{h_1} = 45 \text{ GeV}, f = 100 \text{ GeV}$$



$$m_{h_1} = 50 \text{ GeV}, \beta = 1.5 \times 10^{-4}, m_\chi/f = 0.645$$



- Red line : Relic density
  - Gray & Green : excluded by XENON1T and LEP , respectively
  - Purple : disfavored by diffuse gamma-rays from Fermi-LAT dwarf galaxies
  - Orange : disfavored by Higgs invisible decay
- DM & Nucleon scattering       $h_1 \rightarrow b\bar{b}$

# Summary

- In four-form flux model , a simultaneous relaxation of the Higgs mass and cosmological constant is possible.  
→  $M_{eff}^2(q)$  &  $\Lambda_{eff}(q)$  depend on one flux parameter ‘q’
- We consider pseudo-scalar field to explain the reheating process.
- In this scenario, we introduce Dirac fermion DM with pseudo-scalar coupling.  
→ Flux portals lead to mixing between pseudo-scalar and Higgs  
→ Direct Detection : suppressed  
Indirect Detection : unsuppressed signatures for cosmic rays