

# Bosonic dark matter in coherent state

Eung Jin Chun



**YONSEI**  
UNIVERSITY

---

Based on 2109.07423

# Outline

- Ultralight boson with a huge population is a **cold dark matter** candidate.  
$$\frac{n_\phi}{s} \sim 10^{19} \frac{10^{-20} \text{eV}}{m_\phi}$$
- It can be generated by a misaligned initial amplitude forming a **coherent state** (e.g., QCD axion with instanton potential).
- New idea: **Coherent DM state** driven by **thermal fermions**.

# Bosonic DM in a coherent state

- Coherent state = Eigenstate of annihilation/creation operators

$$\hat{\phi}(x) = \int_k [a_k e^{-ik \cdot x} + a_k^+ e^{+ik \cdot x}]$$

$$a_k |\phi_c\rangle = \phi_k |\phi_c\rangle; a_k^+ |\phi_c\rangle = \phi_k^* |\phi_c\rangle$$

$$\langle a_k \rangle_c = \phi_k; \langle a_k^+ \rangle_c = \phi_k^*$$

- Coherent state = classical wave

$$\hat{\phi}(x) = \int_k [a_k e^{-ik \cdot x} + a_k^+ e^{+ik \cdot x}]$$

Assuming a monochromatic state

$$\langle \hat{\phi}(x) \rangle_c = \phi_0 e^{-ik_0 \cdot x} + \phi_0^+ e^{+ik_0 \cdot x}$$

Energy density:  $\rho_\phi = 2m_\phi^2 |\phi_0|^2$

DM amplitude around us :  $\rho_{DM}^\odot \approx 0.3 \text{GeV/cm}^3 \Rightarrow |\phi_0^\odot| \approx 10^{12} \text{GeV} \frac{10^{-22} \text{eV}}{m_\phi}$

# Misalignment mechanism

- Evolution in the FLRW universe:

$$\langle \hat{\phi}(x) \rangle_T = \phi(t)$$

$$\ddot{\phi}(t) + 3H\dot{\phi}(t) + V'(\phi) = 0$$

- For an approximately free field:

$$\phi''(x) + \frac{3}{2x} \phi'(x) + \phi(x) \approx 0$$

$$x \equiv m_\phi t$$

- Analytic solution

$$\phi(x) = C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}}$$

$$H \gg m_\phi \quad (x \ll 1) : \quad \phi = \phi_i; \dot{\phi} = 0$$

$$H \ll m_\phi \quad (x \gg 1) : \quad \phi \sim \phi_i \frac{\sin(m_\phi t + \frac{\pi}{8})}{(m_\phi t)^{3/4}}$$

$$m_\phi \gg H_{eq} \approx 3 \cdot 10^{-27} \text{ eV}$$

CDM density:  $\rho_{DM}(x_{eq}) \approx 0.48 \text{ eV}^4$      $\rho_\phi(x) \sim \frac{m_\phi^2 \phi_i^2}{x^{3/2}} \Rightarrow \phi_i \sim 0.01 M_p \left( \frac{10^{-20} \text{ eV}}{m_\phi} \right)^{1/4}$

```

ysol =
DSolve[y''[x] + 3/2/x y'[x] + y[x] == 0,
y[x], x][[1]]

{y[x] →  $\frac{\text{BesselJ}\left[\frac{1}{4}, x\right] \mathbb{C}_1}{x^{1/4}} + \frac{\text{BesselY}\left[\frac{1}{4}, x\right] \mathbb{C}_2}{x^{1/4}}$ }

yns =
NDSolve[{y''[x] + 3/2/x y'[x] + y[x] == 0, y[0.01] == 1, (y[x]^2 + y'[x]^2)/2 /. yns;
y'[0.01] == 0}, y, {x, 0.01, 100}][[1]];

Plot[{y[x] /. yns /. x → 10^xx, y'[x] /. yns /. x → 10^xx}, {x, 0.1, 100}, PlotStyle → {Blue, Red}]



Series[y[x] /. ysol, {x, 0, 1}] // FullSimplify

$$-\frac{2^{1/4} \mathbb{C}_2 \Gamma\left[\frac{1}{4}\right]}{\pi \sqrt{x}} + \frac{\mathbb{C}_1 + \mathbb{C}_2}{2^{1/4} \Gamma\left[\frac{5}{4}\right]} + O[x]^{5/4}$$


Series[y[x] /. ysol, {x, Infinity, 1}] // Normal

$$-\frac{\sqrt{\frac{2}{\pi}} \mathbb{C}_2 \cos\left[\frac{\pi}{8} + x\right]}{x^{3/4}} + \frac{\sqrt{\frac{2}{\pi}} \mathbb{C}_1 \sin\left[\frac{\pi}{8} + x\right]}{x^{3/4}}$$


```



# Scalar field in thermal background

- Scalar field interacting with thermal fermions:

$$\mathcal{L}' = y_\phi \hat{\phi} (\bar{f}_R f_L + \bar{f}_L f_R)$$

$$\rightarrow V_{T,\text{eff}}(\phi) = -\frac{g_f}{2\pi^2} T^4 J_F \left( \frac{(m_f + y_\phi \phi)^2}{T^2} \right)$$

Dolan+Jackiw,  
Weinberg, 1974

- Leading thermal effects in cosmological evolution:

$$\ddot{\phi}(t) + 3H\dot{\phi}(t) + (m_\phi^2 + m_T^2)\phi(t) \approx \frac{\partial}{\partial \phi} \langle \mathcal{L}' \rangle_T$$

$$m_T^2 = \frac{g_f}{24} y_\phi^2 T^2, \quad \langle \mathcal{L}' \rangle_T = y_\phi \phi \frac{g_f m_f T^2}{24}$$

$$g_f = 4N_c \quad (2) \text{ for } f = q, l \text{ (v)}$$

Esteban+Salvado, 2101.05804

Batell+Ghalsasi, 2109.04476

# General features I

- Evolution from  $T_{ew} \approx 100$  GeV down to  $T_{eq} \approx 0.8$  eV:

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \left(1 + \frac{x_1}{x}\right) \tilde{\phi}(x) = \frac{x_s}{x}$$

$$x_1 \equiv y_\phi^2 \frac{c_t^2 g_f}{48} \frac{M_P}{m_\phi}, \quad x_s \equiv y_\phi \frac{c_t^2 g_f}{48} \frac{m_f}{m_\phi}$$

(Notation)  $T = c_t \sqrt{M_P/2t}$   $c_t = 1.74/g_*^{1/4}$

$$x = m_\phi t = \frac{c_t^2 m_\phi M_P}{2T^2},$$

$$\tilde{\phi} = \phi/M_P, \quad m_{20} \equiv m_\phi/10^{-20} \text{eV}$$

- Vanishing initial condition at  $x_{ew}$ :  $\phi = 0, \phi' = 0$ .

- Nontrivial evolution from  $T_{ew}$  ( $x_{ew}$ ) to  $T_f = m_f(x_f)$ , and then free evolution from  $T_f(x_f)$  to  $T_{eq}$  ( $x_{eq}$ ) for  $f = q, l$ .

- Evolution from  $T_{ew}$  to  $T_{eq}$  for  $f = \nu$ .

# General features II

- Different types of solutions depending on the location of  $T_1(x_1)$ :

$$x_{ew} \longrightarrow x_f \longrightarrow x_{eq}$$

For  $f = q, l$ :  
i)  $x_1 < x_{ew} < x_f \rightarrow x_{eq}$   
ii)  $x_{ew} < x_f < x_1 \rightarrow x_{eq}$   
iii)  $x_{ew} < x_1 < x_f \rightarrow x_{eq}$

For  $f = \nu$ :  $x_{ew} < x_1 < x_{eq}$

- The resulting DM density is to be described by

$$\rho_\phi(x_{eq}) \approx \frac{c_f^2 x_S^2}{\pi} \frac{m_\phi^2 M_P^2}{x_{eq}^{3/2}}$$

$$\rho_\phi = \rho_{DM} \Rightarrow c_f^2 x_S^2 \approx 2.5 \times 10^{-4} m_{20}^{-1/2}$$

$$x_{ew} \approx 10^{-15} m_{20} \quad x_{eq} \approx 2 \times 10^7 m_{20}$$

$$x_f \approx 5 \times 10^{-4} m_{20} (m_e/m_f)^2$$

$$x_1 \approx 10^{47} y_\phi^2 m_{20}^{-1} \quad x_S \approx 2 \cdot 10^{24} y_\phi m_{20}^{-1} \left( \frac{m_f}{m_e} \right)$$

# Asymptotic solutions in various regimes

$x \gg x_1$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \tilde{\phi}(x) = \frac{x_S}{x}$$

$x \ll x_1$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \frac{x_1}{x} \tilde{\phi}(x) = \frac{x_S}{x}$$

$x > x_f$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

$$\begin{aligned} \tilde{\phi}(x) = & \frac{\pi x_S}{(2x)^{1/4}} \left( \frac{x}{\Gamma(3/4)} J_{1/4}(x) F_{1,2} \left( \frac{1}{2}; \frac{3}{4}, \frac{3}{2}; -\frac{x^2}{4} \right) \right. \\ & - \frac{x^{3/2}}{3\Gamma(5/4)} \left( J_{1/4}(x) - Y_{1/4}(x) \right) F_{1,2} \left( \frac{3}{4}; \frac{5}{4}, \frac{7}{4}; -\frac{x^2}{4} \right) \Big) \\ & + C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}} \end{aligned}$$

$$\tilde{\phi}(x) = \frac{x_S}{x_1} + C_1 \frac{e^{2i\sqrt{xx_1}}}{\sqrt{x}} + iC_2 \frac{e^{-2i\sqrt{xx_1}}}{\sqrt{xx_1}}$$

$$\tilde{\phi}(x) = C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}}$$

$\tilde{\phi}(x)$  at  
 $x \rightarrow 0$  &  $\infty$

Case i)

```
DSolve[y''[x] + 3/2/x y'[x] + y[x] == xS/x, y[x], x][[1]];
Series[y[x] /. %, {x, 0, 1}] // FullSimplify
Series[y[x] /. %%, {x, Infinity, 1}] // Normal // FullSimplify
```

$$-\frac{2^{1/4} c_2 \Gamma\left[\frac{1}{4}\right]}{\pi \sqrt{x}} + \frac{c_1 + c_2}{2^{1/4} \Gamma\left[\frac{5}{4}\right]} + \frac{2 x S x}{3} + O[x]^{5/4}$$

$$\frac{x S}{x} + \frac{-\cos\left[\frac{\pi}{8} + x\right] (2 c_2 + 2^{1/4} \sqrt{\pi} x S \Gamma\left[\frac{3}{4}\right]) + 2 c_1 \sin\left[\frac{\pi}{8} + x\right]}{\sqrt{2 \pi} x^{3/4}}$$

Case iii)

```
DSolve[y''[x] + 3/2/x y'[x] + y[x] == 0, y[x], x][[1]];
Series[y[x] /. %, {x, 0, 1}] // FullSimplify
Series[y[x] /. %%, {x, Infinity, 1}] // Normal
```


$$-\frac{2^{1/4} c_2 \Gamma\left[\frac{1}{4}\right]}{\pi \sqrt{x}} + \frac{c_1 + c_2}{2^{1/4} \Gamma\left[\frac{5}{4}\right]} + O[x]^{5/4}$$

$$-\frac{\sqrt{\frac{2}{\pi}} c_2 \cos\left[\frac{\pi}{8} + x\right]}{x^{3/4}} + \frac{\sqrt{\frac{2}{\pi}} c_1 \sin\left[\frac{\pi}{8} + x\right]}{x^{3/4}}$$


# Case i)

- Behaviors of  $\phi$  &  $\rho_\phi$  with the vanishing initial conditions

```
yns =
NDSolve[{y''[x] + 3/2/x y'[x] + y[x] == 1/x, y[0.01] == 1,
y'[0.01] == 0}, y, {x, 0.01, 100}][[1]];
Plot[{y[x] /. yns /. x → 10^xx, y'[x] /. yns /. x → 10^xx},
{xx, -2, 2}, PlotStyle → {Blue, Red}]
```



```
(y[x]^2 + y'[x]^2) / 2 /. yns;
LogLogPlot[%, % x^(3/2), (100/x)^2 (x/100)^(3/2),
{x, 0.1, 100}, PlotStyle → {Blue, Red, Yellow}]
```




## Case iii)

- Behaviors of  $\phi$  &  $\rho_\phi$  with the initial misalignment.

```

yns =
NDSolve[{y''[x] + 3/2/x y'[x] + y[x] == 0, y[0.01] == 1,
y'[0.01] == 0}, y, {x, 0.01, 100}][[1]];
Plot[{y[x] /. yns /. x → 10^xx, y'[x] /. yns /. x → 10^xx},
{xx, -2, 2}, PlotStyle → {Blue, Red}]


```



```

(y[x]^2 + y'[x]^2) / 2 /. yns;
LogLogPlot[ %, % x^(3/2), 0.4 (100/x)^2 (x/100)^(3/2),
{x, 0.1, 100}, PlotStyle → {Blue, Red, Yellow}]

```



# Scalar DM condensed by $q/l$

i)  $x_1 < x_{ew} < x_f$

$$x < x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

## Solution

$$\tilde{\phi}(x) = \begin{cases} x_S \left( \frac{\pi}{\Gamma(\frac{3}{4})} \frac{J_{1/4}(x)}{(2x)^{1/4}} [G_1(x) - G_1(x_{ew})] - \frac{4\Gamma(\frac{3}{4})}{3} \frac{J_{-1/4}(x)}{(2x)^{1/4}} [G_2(x) - G_2(x_{ew})] \right) \\ C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}} \end{cases}$$

$$G_1(x) = x F_{1,2} \left( \frac{1}{2}; \frac{3}{4}, \frac{3}{2}; -\frac{x^2}{4} \right), \quad G_2(x) = x^{3/2} F_{1,2} \left( \frac{3}{4}; \frac{5}{4}, \frac{7}{4}; -\frac{x^2}{4} \right)$$

$$\frac{C_1}{x_S} = \frac{\pi}{2^{\frac{1}{4}} \Gamma(\frac{3}{4})} G_1(x_f) - \frac{2^{\frac{5}{4}} \Gamma(\frac{3}{4})}{3} G_2(x_f)$$

$$\frac{C_2}{x_S} = \frac{2^{\frac{5}{4}} \Gamma(\frac{3}{4})}{3} G_2(x_f)$$

# Scalar DM condensed by $q/l$

i)  $x_1 < x_{\text{ew}} < x_f$

$$x < x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \frac{x_1}{x} \tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

$$\rho_\phi(x_{eq}) \approx \frac{c_f^2 x_S^2 m_\phi^2 M_P^2}{\pi x_{eq}^{3/2}} \text{ with } c_f^2 = C_1^2 + C_2^2$$

$$c_f^2 = \begin{cases} 4\sqrt{2}\Gamma\left(\frac{5}{4}\right)^2 x_f^2 & \text{for } x_f \ll 1 \\ \frac{\pi\Gamma\left(\frac{3}{4}\right)^2}{2\sqrt{2}} & \text{for } 1 \ll x_f \end{cases}$$

$$y_\phi \approx 3.6 \cdot 10^{-24} \frac{m_{20}^{-\frac{1}{4}}}{N_c} \left(\frac{m_f}{m_e}\right) \text{ with } \frac{550}{N_c^{\frac{2}{5}}} \left(\frac{m_f}{m_e}\right)^{\frac{4}{5}} \ll m_{20} \ll 2 \cdot 10^3 \left(\frac{m_f}{m_e}\right)^2 \text{ for } x_f \ll 1$$

$$y_\phi \approx 3 \cdot 10^{-27} \frac{m_{20}^{\frac{3}{4}}}{N_c} \left(\frac{m_e}{m_f}\right) \text{ with } m_{20} \gg \text{Max} \left[ 2 \frac{10^{16}}{N_c^2} \left(\frac{m_f}{m_e}\right)^4, 2 \cdot 10^3 \left(\frac{m_f}{m_e}\right)^2 \right] \text{ for } 1 \ll x_f$$

$m_\phi \ll 46 \text{ eV (5.7 MeV)} \text{ for } f = e \text{ (b)},$   
 (\*) Neglecting Freeze-In:  $y_\phi \ll 5.3 (1.4) \times 10^{-11}.$

# Scalar DM condensed by $q/l$

ii)  $x_{ew} < x_f < x_1$

$$x < x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \frac{x_1}{x}\tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

## Solution

$$\tilde{\phi}(x) = \begin{cases} \frac{x_S}{x_1} \left( 1 - \sqrt{\frac{x_{ew}}{x}} \cos[2(\sqrt{xx_1} - \sqrt{x_{ew}x_1})] - \frac{1}{2\sqrt{xx_1}} \sin[2(\sqrt{xx_1} - \sqrt{x_{ew}x_1})] \right) \\ C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}} \end{cases}$$

$$\frac{C_1}{x_S} = -\frac{\pi x_f^{\frac{1}{4}}}{4x_1} \left[ 2x_f Y_{\frac{5}{4}}(x_f) - Y_{\frac{1}{4}}(x_f) \cos(2\sqrt{x_1 x_f}) + \sqrt{\frac{x_f}{x_1}} Y_{-\frac{3}{4}}(x_f) \sin(2\sqrt{x_1 x_f}) \right],$$

$$\frac{C_2}{x_S} = \frac{\pi x_f^{\frac{1}{4}}}{4x_1} \left[ 2x_f J_{\frac{5}{4}}(x_f) - J_{\frac{1}{4}}(x_f) \cos(2\sqrt{x_1 x_f}) + \sqrt{\frac{x_f}{x_1}} J_{-\frac{3}{4}}(x_f) \sin(2\sqrt{x_1 x_f}) \right].$$

# Scalar DM condensed by $q/l$

ii)  $x_{ew} < x_f < x_1$

$$x < x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \frac{x_1}{x}\tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

$$\rho_\phi(x_{eq}) \approx \frac{c_f^2 x_S^2}{\pi} \frac{m_\phi^2 M_P^2}{x_{eq}^{3/2}} \text{ with } c_f^2 = C_1^2 + C_2^2$$

$$c_f^2 = \begin{cases} 4\sqrt{2}\Gamma\left(\frac{5}{4}\right)^2 x_f^2 & \text{for } x_f \ll x_1 \ll 1 \\ \frac{\pi x_f^3}{2x_1^2} & \text{for } 1 \ll x_f \ll x_1 \end{cases}$$

$$y_\phi \approx 3.6 \cdot 10^{-24} \frac{m_{20}^{-\frac{1}{4}}}{N_c} \left(\frac{m_f}{m_e}\right) \text{ with } m_{20} \ll 31 N_c^{-\frac{2}{5}} \left(\frac{m_f}{m_e}\right)^{8/5} \text{ for } x_f \ll x_1 \ll 1$$

$$y_\phi \approx 5.3 \cdot 10^{-24} m_{20} \left(\frac{m_e}{m_f}\right)^{\frac{1}{2}} \text{ with } m_{20} \gg 2 \cdot 10^3 \left(\frac{m_f}{m_e}\right)^2 \text{ for } 1 \ll x_f \ll x_1$$

(\*) BBN constraint:  $m_\phi \ll 10^{-6} (2.5 \times 10^{-3}) \text{ eV}$  for  $f = e$  ( $b$ ),

$$y_\phi \ll 5.7 \times 10^{-10} (1.5 \times 10^{-8}).$$

# Scalar DM condensed by $q/l$

iii)  $x_{ew} < x_1 < x_f$

$$x < x_1$$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \frac{x_1}{x} \tilde{\phi}(x) = \frac{x_S}{x}$$

$$x_1 < x < x_f$$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \tilde{\phi}(x) = \frac{x_S}{x}$$

$$x_f < x$$

$$\tilde{\phi}''(x) + \frac{3}{2x} \tilde{\phi}'(x) + \tilde{\phi}(x) = 0$$

(i)  $x_1 \ll x_f \ll 1$ :

$$y_\phi \approx 3.6 \times 10^{-24} \frac{m_{20}^{-\frac{1}{4}}}{N_c} \frac{m_f}{m_e} \quad \text{with} \quad \frac{28}{N_c^{\frac{2}{5}}} \left( \frac{m_f}{m_e} \right)^{\frac{8}{5}} \ll m_{20} \ll \text{Min} \left[ 2 \times 10^3 \left( \frac{m_f}{m_e} \right)^2, \frac{6 \times 10^5}{N_c^{\frac{2}{5}}} \left( \frac{m_f}{m_e} \right)^{\frac{4}{5}} \right]$$

(ii)  $x_1 \ll 1 \ll x_f$ :

$$y_\phi \approx 3 \times 10^{-27} \frac{m_{20}^{\frac{3}{4}}}{N_c} \frac{m_e}{m_f} \quad \text{with} \quad 2 \times 10^3 \left( \frac{m_f}{m_e} \right)^2 \ll m_{20} \ll \text{Min} \left[ 3.2 \times 10^{11} N_c^2 \left( \frac{m_f}{m_e} \right)^4, \frac{2.1 \times 10^{16}}{N_c^2} \left( \frac{m_e}{m_f} \right)^4 \right]$$

(iii)  $1 \ll x_1 \ll x_f$ :

$$y_\phi \approx 1.9 \times 10^{-22} \frac{m_{20}^{\frac{1}{3}}}{N_c^{\frac{1}{6}}} \left( \frac{m_f}{m_e} \right)^{\frac{2}{3}} \quad \text{with} \quad 1.2 \times 10^8 N_c^{\frac{5}{4}} \left( \frac{m_f}{m_e} \right)^{\frac{13}{4}} \ll m_{20} \ll 4.4 \times 10^{11} N_c^2 \left( \frac{m_f}{m_e} \right)^4 \quad \text{for } x_1^5 \ll x_f$$

$$y_\phi \approx 4.5 \times 10^{-28} \frac{m_{20}}{N_c} \left( \frac{m_e}{m_f} \right)^{\frac{3}{2}} \quad \text{with} \quad m_{20} \gg 4.9 \times 10^8 N_c^{\frac{5}{4}} \left( \frac{m_f}{m_e} \right)^{\frac{13}{4}} \quad \text{for } x_f \ll x_1^5.$$

(\*) BBN+Feeze-In:  $y_\phi \ll 5.7 (1.2) \times 10^{-10}$ .

$m_\phi \ll 1.3 \times 10^{-2} (6.1 \times 10^3)$  eV for  $f = e$  (b)

# Scalar DM condensed by $\nu$

- $x_{ew} < x_1 < x_{eq} < x_\nu$

$$x < x_1$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \frac{x_1}{x}\tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_1$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = \frac{x_S}{x}$$

## Solution

$$\tilde{\phi}(x) = \begin{cases} \frac{x_S}{x_1} \left( 1 - \sqrt{\frac{x_{ew}}{x}} \cos[2(\sqrt{xx_1} - \sqrt{x_{ew}x_1})] - \frac{1}{2\sqrt{xx_1}} \sin[2(\sqrt{xx_1} - \sqrt{x_{ew}x_1})] \right) \\ C_1 \frac{J_{1/4}(x)}{x^{1/4}} + C_2 \frac{Y_{1/4}(x)}{x^{1/4}} + \frac{x_S}{(2x)^{1/4}} \left( \frac{\pi}{\Gamma(\frac{3}{4})} J_{1/4}(x) G_1(x) - \frac{4\Gamma(\frac{3}{4})}{3} J_{-1/4}(x) G_2(x) \right) \end{cases}$$

(\*)  $G_{1,2}(x) \rightarrow \text{constant for } x \rightarrow \infty$

$$\frac{C_1}{x_S} = -\frac{\pi}{2^{\frac{1}{4}}\Gamma(\frac{3}{4})} G_1(x_1) + \frac{2^{\frac{5}{4}}\Gamma(\frac{3}{4})}{3} G_2(x_1) - \frac{\pi}{2} \frac{Y_{\frac{1}{4}}(x_1) \sin^2(x_1) - Y_{-\frac{3}{4}}(x_1)[x_1 - \frac{1}{2}\sin(2x_1)]}{x_1^{3/4}}$$

$$\frac{C_2}{x_S} = -\frac{2^{\frac{5}{4}}\Gamma(\frac{3}{4})}{6} G_2(x_1) + \frac{\pi}{2} \frac{J_{\frac{1}{4}}(x_1) \sin^2(x_1) - J_{-\frac{3}{4}}(x_1)[x_1 - \frac{1}{2}\sin(2x_1)]}{x_1^{3/4}}$$

# Scalar DM condensed by $\nu$

- $x_{ew} < x_1 < x_{eq} < x_\nu$

$$x < x_1$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \frac{x_1}{x}\tilde{\phi}(x) = \frac{x_S}{x}$$

$$x > x_1$$

$$\tilde{\phi}''(x) + \frac{3}{2x}\tilde{\phi}'(x) + \tilde{\phi}(x) = \frac{x_S}{x}$$

$$\rho_\phi(x_{eq}) \approx \frac{c_\nu^2 x_S^2}{\pi} \frac{m_\phi^2 M_P^2}{x_{eq}^{3/2}} \text{ with } c_\nu^2 = C_1^2 + C_2^2$$

$$c_\nu^2 = \frac{\pi \Gamma\left(\frac{3}{4}\right)^2}{2\sqrt{2}} + \frac{2^{\frac{3}{4}} \pi^{\frac{3}{2}} C_1}{\Gamma\left(\frac{1}{4}\right) x_S} + \frac{C_1^2 + C_2^2}{x_S^2} \approx 3.34 \text{ for } x_1 \gg 1$$

$$y_\phi \approx 4.2 \cdot 10^{-20} m_{20}^{\frac{3}{4}} \left( \frac{0.05 \text{ eV}}{m_\nu} \right) \text{ with } m_{20} \gg 710 \left( \frac{0.05 \text{ eV}}{m_\nu} \right)^4$$

$$\frac{x_S}{x_1} \sim 4.8 \times 10^{-11} m_{20}^{-3/4}$$

$$\frac{x_1}{x_{eq}} \sim 8.8 m_{20}^{-1/2}$$

$$(*) \text{ BBN: } y_\phi \lesssim 7 \times 10^{-6} \quad m_\phi \lesssim 0.092 \text{ eV} \left( \frac{m_\nu}{0.05 \text{ eV}} \right)^{4/3}$$

# Conclusion

- An ultralight scalar can form a coherent DM state through its tiny coupling to SM fermions.
- Considering the  $\phi qq$  or  $\phi ll$  coupling, it works for wide ranges of  $m_\phi \sim (10^{-22}, 10^6)$ eV and  $y_\phi \sim (10^{-28}, 10^{-8})$ .
- For the  $\phi vv$  coupling, DM genesis requires  $m_\phi \sim (10^{-17}, 10^{-1})$ eV and  $y_\phi \sim (10^{-17}, 10^{-5})$ , and can occur **very late** at around  $T > 10T_{eq}$ .